

6. Diagnostic and Statistical Manual of Mental Disorders (DSM-II), Committee on Nomenclature and Statistics of the American Psychiatric Association. Washington, 1968, p 30
7. Kennedy AC, Linton AL, Eaton JC: Urea levels in cerebrospinal fluid after haemodialysis. *Lancet* 1:410-411, 1962
8. Gilliland KG, Hegstron RM: The effect of hemodialysis on cerebrospinal fluid pressure in uremic dogs. *Trans Amer Soc Artif Intern Organs* 9:44-50, 1963
9. Peterson H de C, Swanson A: Acute encephalopathy occurring during hemodialysis—The reverse urea effect. *Arch Intern Med* 113:877-880, 1964
10. Rosen SM, O'Connor K, Shaldon S: Haemodialysis disequilibrium. *Br Med J* 2:672, 1964
11. Brown H, Maher J, Lapierre L, et al: Clinical problems related to the prolonged artificial maintenance of life by hemo-

- dialysis in chronic renal failure. *Trans Amer Soc Artif Intern Organs* 8:281-291, 1962
12. Cooper AJ: Hypomanic psychosis precipitated by hemodialysis. *Comp Psychiat* 8:168-174, 1967
13. Wright R, Sand S, Livingston G: Psychological stress during hemodialysis for chronic renal failure. *Ann Int Med* 64:611-627, 1966
14. Locke S, Merrill J, Tyler H: Neurologic complication of acute uremia. *Arch Intern Med* 108:519-539, 1961
15. Tyler HR: Neurological complications of dialysis, transplantation, and other forms of treatment in chronic uremia. *Neurol* 15:1081-1088, 1965
16. Gonzalez F, Pabico R, Brown H, et al: Further experience with the use of intermittent hemodialysis in chronic renal failure. *Trans Amer Soc Artif Intern Organs* 9:11-17, 1963

Refer to: Goldstein E, Hoepflich PD: Lymphangiography in diagnosis of fever of unknown origin. *Calif Med* 119:59-61, Nov 1973

Lymphangiography in Diagnosis of Fever of Unknown Origin

ELLIOT GOLDSTEIN, MD
PAUL D. HOEPRICH, MD
Davis

WHEN FEVER PERSISTS and the cause remains unknown despite extensive evaluation, surgical exploration of the abdomen is frequently recommended for diagnosis.¹⁻⁴ This procedure has considerable merit because intra-abdominal processes that may cause fever—for example, lymphomas of the retroperitoneal space^{3,5} are frequently occult. However, there are many extra-abdominal causes for persistent fevers that are quite important (among them tuberculosis, subacute bacterial endocarditis, polyarteritis) and these may also elude early diagnosis. As a result, the decision to operate and the timing of the operation are clinical judgments which are exceedingly difficult. We recently performed lymphangiography as part of the preoperative evaluation of a patient with a persistent fever. Because such use of lymphangiography has been reported infrequently,^{3,6} we report our experience to emphasize the value of this procedure in the early diagnosis of fever of unknown origin.

From the Section of Infectious and Immunologic Diseases, Department of Internal Medicine, School of Medicine, University of California, Davis; and the Sacramento Medical Center.

Submitted April 3, 1973.

Reprint requests to: E. Goldstein, MD, Section of Infectious and Immunologic Diseases, Department of Internal Medicine, University of California, Davis, School of Medicine, Davis, CA 95616.

Report of a Case

A 55-year-old male agronomist was referred to the Sacramento Medical Center in April 1972 for evaluation of fever and fatigue of two months' duration. With the exception of these symptoms the history was unremarkable. On physical examination the patient was observed to be chronically ill-appearing, the temperature was 39.8°C (103.6°F), blood pressure 130/80 mm of mercury and pulse rate 108 per minute. A non-pruritic maculopapular rash was present over the chest and abdomen. There was no palpable enlargement of lymph nodes, liver or spleen.

Laboratory studies showed a hematocrit of 35, a leukocyte count of 5,100 per cu mm with 49 percent polymorphonuclear neutrophils, 32 percent lymphocytes, 18 percent monocytes, and 1 percent eosinophiles. The platelet count was 155,000 per cu mm. The erythrocyte sedimentation rate was 53 mm in one hour (Wintrobe). Results of urine and cerebrospinal fluid examinations were normal. Blood chemistry determinations, including glutamic oxaloacetic transaminase (SGOT), alkaline phosphatase, lactic dehydrogenase, bilirubin, albumin, globulin, urea nitrogen, uric acid and haptoglobin were within normal limits. Antinuclear antibody, rheumatoid factor tests, direct and indirect Coombs tests, the VDRL test and serum protein electrophoresis were negative. The feces was guaiac-negative. An electrocardiogram was normal.

No abnormalities were noted in roentgenological examinations of the chest, skull, esophagus, stomach, subphrenic space, small intestine, large intestine, gall bladder, and common bile duct. An intravenous pyelogram revealed a horseshoe kidney. A radionuclide scan of the liver was normal. Skin tests with coccidioidin, intermediate strength tuberculin, histoplasmin, and mumps antigen were nonreactive. Agglutinins for typhoid O, typhoid




Figure 1.—Lymphangiogram showing enlarged para-aortic lymph nodes which are foamy in appearance and contain filling defects.

H, paratyphoid A, paratyphoid B, *Pasteurella tularensis* and *Brucella abortus* were negative, as were blood smears for malaria, and examination of feces for ova and parasites. Cultures of blood, and urine were sterile. A biopsy of bone marrow was interpreted as showing granulocytic hyperplasia. A liver biopsy (Menghini) showed no abnormality. Chronic inflammation was noted in a biopsy specimen from the maculopapular skin rash. Cultures of the biopsy material did not grow either nonacid-fast or acid-fast bacteria, or fungi.

Lymphangiography on the tenth hospital day disclosed enlarged para-aortic abdominal lymph nodes which were compatible in appearance with malignant lymphoma (Figure 1). Abdominal laparotomy was carried out three days later. Biopsy specimens from the enlarged lymph nodes which lay beneath the horseshoe kidney were diagnostic of Hodgkin's disease. The tumor was also present in a liver biopsy specimen and in the excised spleen.

Postoperatively, the patient continued to have high spiking fevers. He was treated with methyl prednisolone, cyclophosphamide, vincristine, and procarbazine. After transient improvement, leukopenia developed with fewer than 300 white blood

cells per cu mm—all lymphocytes. Despite treatment with gentamicin (1.7 mg per kg of body weight every 8 hours, intravenously) and carbenicillin (400 mg per kg of body weight per day by continuous intravenous infusion) the patient died of pneumonia. *Pseudomonas aeruginosa* and *Escherichia coli* grew on cultures of the fluid.

The significant findings at autopsy were Hodgkin's disease involving the bone marrow, liver, and lymph nodes (pelvic, abdominal thoracic, cervical). There was agranulocytosis of the bone marrow and five-lobed, hemorrhagic bacterial pneumonia caused by *P. aeruginosa* and *E. coli* (susceptible, by *in vitro* testing, to gentamicin and carbenicillin).

Discussion

This case illustrates the value of lymphangiography in evaluating fever of occult origin. The roentgenographic features were sufficiently distinctive to predict that a lymphoma involving the para-aortic lymph nodes would be found at laparotomy. However, additional experience is needed to assess the reliability of the procedure. It is likely that falsely positive results occur due to nonspecific lymphadenitis⁸ or to other neoplasms.⁹ Of equal concern is the possibility of falsely negative results—a possibility that is real but apparently small with lymphoma.^{7,8} Although these potential sources for error exist, the high degree of correlation between positive histologic and radiographic findings—more than 80 percent—suggests that the technique will be diagnostically reliable.

The major hazard from lymphangiography is the rare occurrence of an oil embolus.^{10,11} Absolute or relative contraindications are few; they include parenchymal pulmonary disease, a patent foramen ovale and allergy to iodine.⁸

Lymphangiography is also of value in localizing the site of lymph node involvement. In our patient, the lymphangiogram located grossly diseased nodes beneath the horseshoe kidney enabling the surgeon to select an appropriate site for biopsy.

Summary

Lymphangiography of the retroperitoneal lymphatic system was performed early in the evaluation of a patient with a persistent fever of unknown cause. The roentgenograms demonstrated enlarged para-aortic lymph nodes which were compatible in appearance with a malignant lymphoma. Abdomi-

nal laparotomy confirmed the preoperative diagnosis.

Retroperitoneal lymphangiography is recommended as a useful procedure in the preoperative evaluation of patients with persistent fevers of unknown cause.

REFERENCES

1. Geraci JE, Weed LA, Nichols DR: Fever of obscure origin—The value of abdominal exploration in diagnosis. *JAMA* 169: 1306-1315, 1959
2. Hoeprich PD: Resistance to infection, Chap 4, *In* Hoeprich PD (Ed): *Infectious Diseases*. Harper and Row Publishers, 1972, p 49-56
3. Molavi A, Weinstein L: Persistent perplexing pyrexia—Some comments on etiology and diagnosis. *Med Clin N A* 54:379-396, 1970

4. Howard PH Jr, Hardin WJ: The role of surgery in fever of unknown origin. *Surg Clin N A* 52:397-403, 1972

5. Bennett IL Jr, Petersdorf RG: Chills and fever, Chap 16, *In* Wintrobe MM, Thorn GW, Adams RD, et al (Eds): *Harrison's Principles of Internal Medicine*, 6th Ed. McGraw Hill Book Company, 1970, pp 84-92

6. Constantinides S, McPeak C: Lymphangiography and its applications. *Surgery* 57:515-521, 1965

7. Takahashi M, Abrams HL: The accuracy of lymphangiographic diagnosis in malignant lymphoma. *Radiology* 89:448-460, 1967

8. Lee BJ, Martin RS: Indications for lymphangiography in lymphomas and carcinomas. *Med Clin N A* 50:675-688, 1966

9. Lee BJ, Nelson JH, Schwarz G: Evaluation of lymphangiography, inferior venacavography and intravenous pyelography in the clinical staging and management of Hodgkin's disease and lymphosarcoma. *N Engl J Med* 281:327-337, 1964

10. Bron KM, Baum S, Abrams HL: Oil embolism in lymphangiography. *Radiology* 80:194-202, 1963

11. Fuchs WA: Complications of lymphangiography with oily contrast media. *Acta Radiol* 57:427-432, 1962

Scientific Exhibits

103rd Annual Session of the
CALIFORNIA MEDICAL ASSOCIATION
and Western Scientific Assembly

MARCH 2-6, 1974

HILTON HOTEL, SAN FRANCISCO

Send application form for space

SCIENTIFIC AND ORGANIZATIONAL EXHIBITS
1974 Annual Session

To _____

Address _____

City _____ State _____ Zip _____

Mail to CMA Annual Session, 693 Sutter, San Francisco 94102